



β-Carotene Capsule Analyzed with HPLC – AppNote

Separation from Matrix Peaks

Click [HERE](#) for Column Ordering Information.

Beta-carotene may be taken as a dietary supplement in capsule form. In this case, a wide variety of Matrix Peaks were observed in the chromatographic data. It is possible that some of these peaks are various isomers of all-trans β-carotene or other similar carotenes. In any case, resolution was obtained from the other Matrix Peaks, which allows for accurate quantitation of β-carotene in the capsule.



Peak:

β-Carotene

Method Conditions

Column: Cogent Phenyl Hydride™, 4μm, 100Å

Catalog No.: 69020-7.5P

Dimensions: 4.6 x 75 mm

Mobile Phase:

A: DI Water / 0.1% Formic Acid (v/v)

B: Acetonitrile / 0.1% Formic Acid (v/v)

Gradient:

T
i
m
e
(
m
i
n
u
t
e
s
)
7
0
7
0
9
0
6
0
7
0

Post Time: 3 minutes

Injection vol.: 10µL

Flow rate: 1.0 mL / minute

Detection: UV @ 452 nm

Sample Preparation: A Beta-carotene capsule was opened and the contents were transferred to a 25mL volumetric flask containing a portion of Methanol. The solution was sonicated 15 minutes and diluted to mark with Methanol. After mixing, a portion was filtered with a 0.45µm Nylon Syringe Filter (MicroSolv Tech Corp.).

t₀: 0.9 minutes

Note: Beta-carotene is found in many fruits and vegetables. It is responsible for the orange color in carrots, pumpkins, sweet potatoes, and others. In terms of nutrition, Beta-carotene is a metabolic precursor to Vitamin A.



Attachment

No 269 B-Carotene Capsule pdf 0.3 Mb [Download File](#)

Printed from the Chrom Resource Center

MicroSolv Technology Corporation

9158 Industrial Blvd. NE, Leland, NC 28451

tel. (732) 380-8900, fax (910) 769-9435

Email: customers@mtc-usa.com

Website: www.mtc-usa.com

Date: 05-12-2024