


# Oleuropein in Olive Leaves Extract Analyzed with HPLC - AppNote

## **Separation from Matrix Peaks**

The main peak (Oleuropein) was observed to be well Separated from matrix components in this HPLC based approach. This Method is useful for analysts that are not using LCMS and therefore need good Chromatographic Separation in order to perform accurate Quantitation. The matrix Peaks do not co-elute with the main Peak, which is required for UV-based Methods. The Method is highly reproducible (5-run overlay shown in the Figure below).



### **Method Conditions**

Column: Cogent Bidentate C18 2.o™, 2.2μm, 120Å

**Catalog No.**: 40218-05P-2 **Dimensions**: 2.1 x 50mm

**Mobile Phase:** 



A: DI Water with 0.1% Formic Acid (v/v)

B: Acetonitrile with 0.1% Formic Acid (v/v)

#### Gradient:

Time (minutes)	%B
0	5
3	5
19	25
20	25
21	5

Post Time: 5 minutes
Injection vol.: 1µL

Flow rate: 0.3mL / minutes

Detection: UV @ 254nm

 $\textbf{Sample Preparation}: 10.0 \text{mg Olive Leaf Extract was added to 1mL volumetric flask. It was then diluted to mark with 50:50 Solvent A / Solvent B diluent. Then it was filtered with a <math>0.45 \mu m$  Nylon Syringe Filter (MicroSolv Technology).

Corp.).

**to**: 0.6 minutes

**Note**: Olive Leaves are food byproducts (after pruning of olive trees) which are full of bioactive compounds. These compounds are potent polyphenols, which show antibacterial, antiviral, anti-cancer, anti-inflammatory, and antioxidant activities. Different extraction procedures are used for selective extraction of polyphenols from olive leaves. An analytical method to monitor and evaluate the resulting extract is needed.



#### Attachment

No 286 Oleuropein in Olive Leaves Extract Analyzed with HPLC pdf 0.3 Mb Download File

Printed from the Chrom Resource Center

MicroSolv Technology Corporation

9158 Industrial Blvd. NE, Leland, NC 28451 tel. (732) 380-8900, fax (910) 769-9435

Email: customers@mtc-usa.com Website: www.mtc-usa.com

Date: 05-11-2024