

## Using triethylamine TEA in Aqueous Normal Phase ANP methods for simple sugars – Tips & Suggestions

**Triethylamine TEA is an organic base and sometimes used in some HPLC methods to adjust the mobile phase pH.**

Get improved peak shape and selectivity when separating **simple sugars** with TEA and Cogent TYPE-C™ columns as long as the concentration of TEA does not exceed 0.1%. This improved peak shape is thought to be due at least partly to an ion-pairing mechanism of the TEA on the surface of the stationary phase material.

Alternatively, ammonium acetate, a near-neutral pH is obtained with this additive, so it is a popular choice for keeping most organic acids ionized, which is generally preferred for ANP methods. In addition, it is LC-MS compatible, in contrast to a near neutral pH phosphate buffer.

*Our research has indicated that TEA may be difficult to remove from columns once introduced therefore if TEA is to be used with a column, the column should be dedicated to methods with TEA. The same research has shown though that TEA can be used to obtain advantageous changes in chromatography in under certain conditions.*



Printed from the Chrom Resource Center

**MicroSolv Technology Corporation**

9158 Industrial Blvd. NE, Leland, NC 28451

tel. (732) 380-8900, fax (910) 769-9435

Email: [customers@mtc-usa.com](mailto:customers@mtc-usa.com)

Website: [www.mtc-usa.com](http://www.mtc-usa.com)

Date: 04-05-2024