



## Furazolidone Analyzed with HPLC - AppNote

### API and Related Compounds Separation

Furazolidone was Separated from related compounds in a commercially available pharmaceutical preparation. Excellent Peak shape and Selectivity from the related compounds was achieved with this Robust Method making it a great choice for QC applications even when using aggressive Mobile Phases that include acids. Molecular weight of Furazolidone is 225.16.



**Peak:**

Furazolidone

### Method Conditions

**Column:** Cogent Bidentate C18™, 4µm, 100Å

**Catalog No.:** 40018-15P

# MICROSOLV

**Dimensions:** 4.6 x 150mm

**Mobile Phase:** 12% Acetonitrile / 88% DI Water / 1% Acetic Acid

**Temperature:** 30°C

**Injection vol.:** 20µL

**Flow rate:** 2mL / minute

**Detection:** UV @ 365nm

**Note:** *This antibiotic drug is used to treat diarrhea and enteritis caused by bacteria or protozoan infection. Also used as a veterinary medicine for salmonids and Myxobolus cerebralis infection.*



## Attachment

**No 73 Furazolidone Analyzed with HPLC pdf** 0.2 Mb [Download File](#)

Printed from the Chrom Resource Center

**MicroSolv Technology Corporation**

9158 Industrial Blvd. NE, Leland, NC 28451

tel. (732) 380-8900, fax (910) 769-9435

Email: [customers@mtc-usa.com](mailto:customers@mtc-usa.com)

Website: [www.mtc-usa.com](http://www.mtc-usa.com)

Date: 07-03-2024