

## Propantheline Bromide HPLC Methods- AppNote


### Orthogonal Methods without Ion-Pairing Agents


Click [HERE](#) for Column Ordering Information.

In the USP assay method for Propantheline Bromide tablets, Sodium Dodecyl Sulfate (SDS) is used as an Ion-Pairing Agent in the Mobile Phase. The long alkyl chain of this additive makes it particularly difficult and time consuming to load and remove from the HPLC column.

Is this method, not only is the need for Ion Pairing Agents eliminated but the assay can be performed in either the Reversed Phase (Figure A) or Aqueous Normal Phase mode (Figure B).

The Mobile Phase solvents can be used for many methods using these columns. Both methods illustrate good repeatability, with each run from the five-run overlays shown in a different color.





**Peak:**

Propantheline Bromide

**Method Conditions**

**Columns:**

Fig. A: Cogent Phenyl Hydride™, 4μm, 100Å

Fig. B: Cogent Diamond Hydride™, 4μm, 100Å

**Catalog No.:**

Fig. A: 69020-7.5P

Fig. B: 70000-7.5P

**Dimensions:**

Fig. A: 4.6 x 75 mm

Fig. B: 4.6 x 75 mm

**Mobile Phase:**

A: DI Water / 0.1% Formic Acid

B: 97% Acetonitrile / 3% DI Water / 0.1% Formic Acid

**Gradient:**

Figure A.

Time (minutes)	%B
0	10
4	50
5	10

Figure B.

Time (minutes)	%B
0	80
4	70
5	80



**Temperature:**

Fig. A: 35°C

Fig. B: 25°C

**Injection vol.:** 10µL

**Flow rate:** 1.0mL / minute

**Detection:** UV @ 254nm

**Sample:**

**Stock Solution:** 1.0mg Propantheline Bromide Diluted with 1 mL of 50:50 A:B

**Working Solution:** 100µL stock diluted with 900µL 50:50 Solvent A: Solvent B

**Note:** Propantheline Bromide is a Muscarinic Acetylcholine receptor antagonist which is used for the treatment of ulcers, cramps and spasms of the digestive system, and hyperhidrosis.



**Attachment**

**No 132 Orthogonal Methods for Propantheline Bromide pdf** 0.3 Mb [Download File](#)