


Analysis of Nucleotides - AppNote

ATP & AMP Separated in Aqueous Normal Phase (ANP)

The chromatogram below shows an example of the separation possible with this Method for three Adenosine analytes using an Aqueous Normal Phase (ANP) Gradient.



Peaks:

1. Adenosine-3',5'-Cyclic Monophosphate
2. Adenosine 5'-Monophosphate (AMP)
3. Adenosine 5'-Triphosphate (ATP)

Method Conditions

Column: Cogent Diamond Hydride™, 4µm, 100Å

Catalog No.: 70000-10P-2

Dimensions: 2.1 x 100mm

Mobile Phase:

A: DI Water / 0.1% Ammonium Formate

B: 90% Acetonitrile / 10% DI Water / 0.1% Ammonium Formate

Gradient:

Time (minutes)	%B
0	95
10	70

Post Time: 5 minutes

Injection vol.: 2 μ L

Flow rate: 0.3mL / minutes

Detection: UV @ 254nm

Sample Preparation: 0.3mg of each Nucleotide in 50% Acetonitrile / DI Water +12% Ammonia

Notes: Nucleotides are important Phosphate-containing compounds that are found in living cells and are associated with a broad array of metabolic and biological processes. They have significant roles in the synthesis of DNA and RNA, are involved in signal transduction pathways, function as coenzymes in biosynthetic pathways and serve as energy reservoirs in biological systems.



Attachment

No 95 Analysis of Nucleotides pdf 0.1 Mb [Download File](#)