



4'-Aminoacetophenone Analysis with HPLC - AppNote

4'-Aminoacetophenone Retained in Acidic Conditions

This Method investigated 4'-Acetaminophenone Retention and Analysis. Adequate Retention and Sensitivity was the result and could be useful in Analysis of the Metabolites of this class of Anti-tumor Agents in body fluids (Plasma or Urine) during or after Chemotherapy.



Peak:

4'-Aminoacetophenone

Method Conditions

Column: Cogent Bidentate C18™, 4µm, 100Å

Catalog No.: 40018-75P

Dimensions: 4.6 x 75mm

Mobile Phase:

Isocratic: 80:20 Solvent A / Solvent B

A: DI Water / 0.2% Acetic Acid

B: Acetonitrile / 0.2% Acetic Acid

Injection vol.: 2µL

Flow rate: 1.0mL / minute

Detection: UV @ 324nm

Sample Preparation: 1mg of the Compound dissolved in 1mL of 50:50 Solvent A / Solvent B solution.

Sample for Injection diluted 1:15 with 100% Solvent A.

Note: 4'-Aminoacetophenone (arylamine) is one of the metabolites of 1-(4-Acetylphenyl)-3,3-Dimethyltriazene, which is an anti-tumour Triazene. 4'-Acetaminophenone is also frequently used as an internal standard in analysis of Mitomycin C.



Attachment

No 66 4'-Aminoacetophenone Analysis with HPLC pdf 0.2 Mb [Download File](#)

Printed from the Chrom Resource Center

MicroSolv Technology Corporation

9158 Industrial Blvd. NE, Leland, NC 28451

tel. (732) 380-8900, fax (910) 769-9435

Email: customers@mtc-usa.com

Website: www.mtc-usa.com

Date: 07-22-2024