



L-a-phenylglycine Antibiotic Analysis with HPLC – AppNote


Comparison using Cogent Silica-C™ and UDC-Cholesterol™ HPLC Columns

In this Method, a non-modified, silica-hydride HPLC Column, Cogent Silica-C, produces good Retention of the important amino acid L(+)- α -Phenylglycine with an Isocratic Mobile Phase. In addition the Retention Times are extremely Reproducible (%RSD 0.2).

The same Method using a Cogent UDC-Cholesterol Column (UDC), silica-hydride with an 11 Carbon Hydrocarbon and a Cholesterol molecule bonded to it, highlights the strong Retention on the Silica-C 4.6 x 75mm Column compared with the longer Cholesterol 4.6 x 150mm Column for this compound.

Both Chromatograms use the same Mobile Phase in the ANP or Aqueous Normal Phase mode of HPLC.





Method Conditions

Columns:

Cogent Silica-C™, 4μm, 100Å

Cogent UDC-Cholesterol, 4μm, 100Å

Catalog Nos.:

40000-75P

[69069-15P](#)

Dimensions:

4.6 x 75mm

4.6 x 150mm

Mobile Phase: A: 80% Acetonitrile / 19.5% DI Water / 0.5% Formic Acid (*pre-mixed*)

Injection vol.: 5μL

Flow rate: 1mL / minute

Detection: UV @ 254nm

Sample Preparation: L-(+)-α-Phenylglycine 0.3mg / mL dissolved in the Mobile Phase

t₀: 0.89 minutes

Note: Phenylglycine is a synthetic amino acid used in manufacturing Lactam antibiotics, such as semi synthetic Cephalosporins and Penicillins.



Attachment

No 46 L-a-phenylglycine Antibiotic Analysis with HPLC pdf 0.3 Mb [Download File](#)

Printed from the Chrom Resource Center

MicroSolv Technology Corporation

9158 Industrial Blvd. NE, Leland, NC 28451

tel. (732) 380-8900, fax (910) 769-9435

Email: customers@mtc-usa.com

Website: www.mtc-usa.com

Date: 07-03-2024