
MICROS

Cancer Prodrug Analyzed with HPLC – AppNote

Separation Achieved with Shape Selectivity and Hydrophobic Interaction

This Method provides an example of Reversed Phase Separation containing a number of related compounds. The properties of the solutes are listed in the Table below.

When comparing the properties of Compounds 2 and 3, it is interesting to note that 2 is more hydrophobic than 3 but it's Retention is shorter. The Method uses Shape Selectivity as well as Hydrophobic Interaction which may explain why the less hydrophobic compound is retained longer. In general, the more polar compounds elute earlier than the more hydrophobic species. Compound 7 has no polar functionalities at all so it is the most retained under Reversed Phase conditions.



5. Saccharin

- 6. Quinolinedione-R2
 - 7. Benzopyran

Copyright 2024, All Rights Apply **MicroSolv Technology Corporation** 9158 Industrial Blvd. NE, Leland, NC 28451 tel. (732) 380-8900, fax (910) 769-9435 Email: customers@mtc-usa.com Website: www.mtc-usa.com

MICROS

Method Conditions

Column: Cogent UDC Cholesterol[™], 4µm, 100Å Catalog No.: 69069-7.5P Dimensions: 4.6 x 75mm Mobile Phase: 60:40 Acetonitrile / DI Water + 0.5% Perchloric Acid Injection vol.: 5µL Flow rate: 1mL / minute Detection: UV @ 270nm Sample Preparation: 1mg / mL of each in Acetonitrile + DI Water

Table:

Solute	Mol Wt.	В рКа	Log P
Cytidine-R1	397	3.73	1.54
Cytidine-R2	454	3.73	3.66
Cytidine-R3	425	3.73	2.66
Quinolinedione-R1	536	-	3.46
Saccharin	536	5.36	2.96
Quinolinedione-R2	520	-	2.74
Benzopyran	396	-	2.80



Attachment

Cancer Prodrug Analyzed with HPLC pdf 0.2 Mb Download File

Printed from the Chrom Resource Center Copyright 2024, All Rights Apply **MicroSolv Technology Corporation** 9158 Industrial Blvd. NE, Leland, NC 28451 tel. (732) 380-8900, fax (910) 769-9435 Email: customers@mtc-usa.com Website: www.mtc-usa.com