MICROS

Polycyclic Aromatic Hydrocarbons Analyzed with HPLC – AppNote

Analysis of PAH by Reversed Phase HPLC

Polycyclic Aromatic Hydrocarbons (*PAH*) determination in soil, food, air, body fluids etc. has been a topic of interest for routine Quality Control or Screening Analyses. This Method offers good Resolution, reproducible Retention Time and Peak Shape for these compounds. In addition to the main components, small peaks due to impurities or decomposition products are well Resolved in the easy, isocratic Method.



Printed from the Chrom Resource Center Copyright 2024, All Rights Apply **MicroSolv Technology Corporation** 9158 Industrial Blvd. NE, Leland, NC 28451 tel. (732) 380-8900, fax (910) 769-9435 Email: customers@mtc-usa.com Website: www.mtc-usa.com



Peaks:

- 1. Naphthalene
- 2. Phenanthrene

3. Guaifenesin

4. Pyrene

I. Impurities or Decomposition Product

Method Conditions

Column: Cogent Bidentate C18[™], 4µm, 100Å Catalog No.: 40018-75P Dimensions: 4.6 x 75mm Mobile Phase: 70:30 Acetonitrile / DI Water Injection vol.: 1µL Flow rate: 0.5mL / minute Detection: UV @ 254nm Sample Preparation: 1mg of each sample was dissolved in 1 mL of the Mobile Phase.

Note: A Polycyclic Aromatic Hydrocarbon is a Hydrocarbon Compound containing only Carbon and Hydrogen—that is composed of multiple Aromatic Rings.



Attachment

No 29 Polycyclic Aromatic Hydrocarbons Analyzed with HPLC pdf 0.2 Mb Download File

Printed from the Chrom Resource Center Copyright 2024, All Rights Apply **MicroSolv Technology Corporation** 9158 Industrial Blvd. NE, Leland, NC 28451 tel. (732) 380-8900, fax (910) 769-9435 Email: customers@mtc-usa.com Website: www.mtc-usa.com