





Acrylamide HPLC Analyzed with HPLC - AppNote

Easy & Precise Retention of a Very Polar Compound

Acrylamide can be difficult to retain with conventional Reversed Phase Methods due to its polar nature. With this Method however, Retention is readily achievable using a simple, Isocratic Mobile Phase. The overlay of two Chromatograms using two different lots of HPLC Columns in the *Figure* below illustrates the Reproducibility and Robustness of this Method.





Peak:

Acrylamide

Method Conditions

Column: Cogent Diamond Hydride™, 4µm, 100Å

Catalog No.: 70000-7.5P

Dimensions: 4.6 x 75mm

Mobile Phase: Acetonitrile with 0.1% Formic Acid

Injection vol.: 1µL

Flow rate: 1.0mL / minute

Detection: UV @ 205nm

Sample Preparation: 0.1mg / mL Acrylamide in Mobile Phase as the Diluent.

t₀: 1.0 minute

***Note:** Acrylamide is a monomer used to synthesize polyacrylamides. It was reported to be present in certain food products in 2002. This has been cause for concern as the monomer form is a known carcinogen and neurotoxin. As such, quantitation of this analyte is of importance in a variety of fields.*



Attachment

No 199 Acrylamide Analyzed with HPLC pdf 0.3 Mb [Download File](#)