



Ranitidine Tablet Analyzed by HPLC - AppNote

Excellent Peak Shape of API From an OTC Tablet

Ranitidine has several amine functional groups that can contribute to significant tailing with Reversed Phase Methods. Data shown below uses an extract from a tablet formulation, illustrating how good Peak shape can be obtained. In addition, three Column lots were used to demonstrate the robustness and precision of this Method.





Peaks:

1. Ranitidine
2. Matrix Component

Method Conditions

Column: Cogent Diamond Hydride™, 4μm, 100Å

Catalog No.: 70000-7.5P

Dimensions: 4.6 x 75mm

Mobile Phase:

A: DI Water / 0.1% Trifluoroacetic acid (v/v)

B: Acetonitrile / 0.1% Trifluoroacetic acid (v/v)

Gradient:

Time (minutes)	%B
0	95
1	95
6	50
7	95

Post Time: 3 minutes



Injection vol.: 2 μ L

Flow rate: 1.0mL / minute

Detection: UV @ 313nm

Sample Preparation: 150mg strength Ranitidine HCl tablet was ground and weighed in a 50mL volumetric flask. A portion of 50:50 Solvent A / Solvent B diluent was added and the flask was sonicated 10 minutes. It was then diluted to mark and filtered with a 0.45 μ m Nylon Syringe Filter (MicroSolv Tech Corp.). The filtrate was diluted 1:100 for injections.

t₀: 0.9 minutes

Note: Ranitidine is an acid reducer that is available over-the-counter. It works as a histamine H₂-receptor antagonist, in contrast to other acid reducers such as omeprazole which are proton pump inhibitors. It is sold under the trade name Zantac®.



Attachment

No 217 Ranitidine Tablet Analyzed by HPLC pdf 0.4 Mb [Download File](#)