



Famotidine Tablet Analyzed with HPLC – AppNote

API Separation from Matrix Component

This Method for Analysis of Famotidine Tablets is easy to perform and produces a Symmetrical Peak Shape for the API. This compound has numerous amines which can be problematic in terms of Peak Shape with conventional Columns. Separation from a component from the tablet extract matrix is obtained as well, illustrating specificity of the Method.

Reproducibility is shown by the overlay of runs from two different Column lots.



Peaks:

1. Matrix Component
2. Famotidine

Method Conditions

Column: Cogent Diamond Hydride™, 4µm, 100Å

Catalog No.: 70000-7.5P

Dimensions: 4.6 x 75mm

Mobile Phase:

A: DI Water with 0.1% Trifluoroacetic Acid (TFA) v/v

B: Acetonitrile with 0.1% Trifluoroacetic Acid (TFA) v/v

Gradient:

Time (minutes)	%B
0	95
2	95
6	50
7	95

Post Time: 3 minutes

Injection vol.: 1µL

Flow rate: 1.0mL / minute

Detection: UV @ 265nm

Sample Preparation: 10mg strength Famotidine tablet was ground and added to a 25mL volumetric flask. A portion of 50:50 Solvent A / Solvent B diluent was added and the flask was sonicated 10 minutes. It was then diluted to mark and filtered with a 0.45µm Nylon Syringe Filter (MicroSolv Tech Corp.).

t₀: 0.9 minutes



Attachment

No 221 Famotidine Tablet Analyzed with HPLC pdf 0.4 Mb [Download File](#)

Printed from the Chrom Resource Center

MicroSolv Technology Corporation

9158 Industrial Blvd. NE, Leland, NC 28451

tel. (732) 380-8900, fax (910) 769-9435

Email: customers@mtc-usa.com

Website: www.mtc-usa.com

Date: 04-05-2024