



## Tapentadol HCl Tablet Analyzed by HPLC - AppNote

### Assay Method for the Analgesic Nucynta

Tapentadol can be a problematic compound for HPLC analysis due to the Amine functional group. Tertiary amines are often particularly difficult to obtain a good peak shape using Reversed Phase methods. Peak tailing has been reported in several published papers in the literature.

With this Method, a sharp Peak is obtained due to the unique retention mode. Data from two column lots shown in the figure illustrates the reproducibility of the Method and its robustness.





**Peak:**

Tapentadol HCl

### Method Conditions

**Column:** Cogent Diamond Hydride™, 4µm, 100Å

**Catalog No.:** 70000-7.5P

**Dimensions:** 4.6 x 75mm

**Mobile Phase:**

A: DI Water / 0.1% Formic Acid (v/v)

B: Acetonitrile / 0.1% Formic Acid (v/v)

**Gradient:**

Time (minutes)	%B
0	95
1	95
6	40
7	95

**Post Time:** 3 minutes

**Injection vol.:** 1µL

**Flow rate:** 1.0mL / minute

**Detection:** UV @ 271nm

**Sample Preparation:** 75mg strength Nucynta® tablet was ground and weighed in a 25mL volumetric flask. A portion of 50:50 Solvent A / Solvent B diluent was added and the flask was sonicated 10 minutes. It was then diluted to mark and filtered with a 0.45µm Nylon Syringe Filter (MicroSolv Tech Corp.). The filtrate was diluted 1:5 for HPLC injections.

**t<sub>0</sub>:** 0.9 minutes

---

**Note:** Tapentadol is an analgesic compound used to treat moderate to severe pain. Its efficacy is due to two modes of action: one is an agonist of the  $\mu$ -opioid receptor and another as a norepinephrine reuptake inhibitor.



## Attachment

**No 234 Nucynta Tapentadol HCl Tablet Analyzed by HPLC pdf** 0.3 Mb [Download File](#)