



A: DI Water / 0.1% Formic Acid (v/v)

B: Acetonitrile / 0.1% Formic Acid (v/v)

**Gradient:**

T  
i  
m  
e  
(  
m  
i  
n  
u  
t  
e  
s  
)  
7  
0  
7  
1  
0  
9  
0  
6  
0  
7  
0

**Post Time:** 3 minutes

**Injection vol.:** 10 $\mu$ L

**Flow rate:** 1.0 mL / minute

**Detection:** UV @ 452 nm

**Sample Preparation:** A Beta-carotene capsule was opened and the contents were transferred to a 25mL volumetric flask containing a portion of Methanol. The solution was sonicated 15 minutes and diluted to mark with Methanol. After mixing, a portion was filtered with a 0.45 $\mu$ m Nylon Syringe Filter (MicroSolv Tech Corp.).

**t<sub>0</sub>:** 0.9 minutes

**Note:** Beta-carotene is found in many fruits and vegetables. It is responsible for the orange color in carrots, pumpkins, sweet potatoes, and others. In terms of nutrition, Beta-carotene is a metabolic precursor to Vitamin A.



**Attachment**

No 269 B-Carotene Capsule pdf 0.3 Mb [Download File](#)

Printed from the Chrom Resource Center  
**MicroSolv Technology Corporation**  
9158 Industrial Blvd. NE, Leland, NC 28451  
tel. (732) 380-8900, fax (910) 769-9435  
Email: [customers@mtc-usa.com](mailto:customers@mtc-usa.com)  
Website: [www.mtc-usa.com](http://www.mtc-usa.com)  
Date: 07-22-2024