


Peonidin-3-Glucoside Analyzed by LCMS - AppNote

LC-MS method for Anthocyanin- Fruit or Vegetable Extracts

Click [HERE](#) for Column Ordering Information.

In this analysis, the presence of Peonidin-3-O-Glucoside was confirmed in two out of four extracts. The peaks were symmetrical and retained beyond the dead volume.

Due to the MS detection and comparison of the retention time, it was possible to find the peak for Peonidin-3-O-Glucoside among many peaks with similar m/z values present in the A1 extract. The method after validation can be used for quality control of commercial fruit extracts as well in studies of bioactivities of this important class of compounds.



Peak:

Peonidin 3-O-glucoside 463 m/z [M+]

Method Conditions

Column: Cogent Phenyl Hydride™, 4µm, 100Å

Catalog No.: [69020-05P-2](#)

Dimensions: 2.1 x 50 mm

Mobile Phase:

A: DI Water / 0.1% Formic Acid (v/v)

B: Acetonitrile / 0.1% Formic Acid (v/v)

Gradient:

Time
min
%
B
min
ul
e
s
)
1
0
5
8
4
0

8
6
0
1
7
5

Post Time: 3 minutes

Injection vol.: 1 μ L

Flow rate: 0.4 mL/minute

Detection: ESI - POS - Perkin Elmer, Flexar SQ 300 Mass Spectrometer

Sample Preparation: Four proprietary fruit or vegetable extracts were analyzed. Samples were marked A1 to A4.

t₀: 0.4 minutes

Note: Anthocyanins are plant pigments and have recently been used as natural colorants in food products instead of synthetic dyes used in the food industry, which have been suspected to cause adverse neurological effects. The rapidly growing food and beverage market demands more and more of plant pigments. Also the main secondary metabolites in black rice are Anthocyanins, mainly Cyanidin-3-Oglucoside and Peonidin-3-O-Glucoside.



Attachment

No 296 Peonidin-3-Glucoside pdf 0.3 Mb [Download File](#)