



B Vitamins Analysis with HPLC - AppNote

Separation of Five Highly Polar Vitamins

The B Vitamins are all highly polar and often ion pair agents are needed in Reverse Phase in order to retain them. However, these agents are incompatible with LCMS and therefore limit the applications of these methods. In this method using the Diamond Hydride Column, good separation is obtained with MS-compatible mobile phase solvents.



Peaks:

1. Niacinamide
2. Riboflavin

3. Pyridoxine
4. Folic Acid
5. Cyanocobalamin

Method Conditions

Column: Cogent Diamond Hydride™, 4µm, 100Å

Catalog No.: 70000-7.5P

Dimensions: 4.6 x 75mm

Mobile Phase:

A: DI Water / 10 mM Ammonium Formate

B: 95% Acetonitrile / 5% Solvent A

Gradient:

Time (minutes)	%B
0	100
2	100
9	50
10	100

Post Time: 2 minutes

Flow rate: 1.0 mL/minute

Detection: UV @ 266 nm

Injection vol.: 2µL

Sample Preparation: Mix of 0.1 mg/L Niacinamide, 0.01 mg /mL Riboflavin, 0.3 mg / mL Pyridoxine, 0.05 mg / mL Folic Acid, 1.0 mg / mL Cyanocobalamin in 50% Solvent A/ 50% Solvent B diluent. Peak identities were confirmed by individual standards.

t₀: 0.9 minutes

Note: The word "Vitamin" was originally spelled "vitamine" when it was first coined by biochemist Casimir Funk. It was derived from the words "vital" and "amine" because it was believed at the time that all Vitamins were chemical Amines. The "e" was dropped from the word when it was discovered that this is not the case.



Attachment

No 162 B Vitamins Analysis pdf 0.6 Mb [Download File](#)

Printed from the Chrom Resource Center
MicroSolv Technology Corporation
9158 Industrial Blvd. NE, Leland, NC 28451
tel. (732) 380-8900, fax (910) 769-9435
Email: customers@mtc-usa.com
Website: www.mtc-usa.com
Date: 07-22-2024