

# MICROSOLV

## **Separation of Isobaric Drugs**

Morphine and Hydromorphone are Isobaric Drugs (the same m/z value) that can be separated using the Cogent Bidentate C18 2.0<sup>™</sup> Column. Multiple samples (n=5) were prepared and analyzed. The results showed great reproducibility (RSDs < 3%). The detection limits were 50 to 500 times lower than that required by immunoassays.





Peaks:

1. Morphine 286.1438 m/z [M+H]+

2. Hydromorphone 286.1438 m/z [M+H]+

## **Method Conditions**

Column: Cogent Bidentate C18 2.0<sup>™</sup>, 2.2µm, 120Å

#### Catalog No.: 40218-05P-2

Dimensions: 2.1 x 50 mm

### **Mobile Phase:**

A: DI Water / 0.1% Formic Acid (v/v)

B: 50% Acetonitrile / 50% Methanol / 0.1% Formic Acid (v/v)

### Gradient:

Time ( <i>minutes</i> )	%B
0	5
4	50

Printed from the Chrom Resource Center Copyright 2024, All Rights Apply **MicroSolv Technology Corporation** 9158 Industrial Blvd. NE, Leland, NC 28451 tel. (732) 380-8900, fax (910) 769-9435 Email: customers@mtc-usa.com Website: www.mtc-usa.com

# MICROSOLV

5	90
6	90
7	5

Post Time: 3 minutes Injection vol.: 1µL Flow rate: 0.4mL/minute Detection: ESI – POS – Agilent 6210 MSD TOF Mass Spectrometer Sample: The drugs were spiked into urine at a level of 50 ng/mL. Ex

Sample: The drugs were spiked into urine at a level of 50 ng/mL. Extraction method: Spiked urine sample was loaded into SPE cartridge I (Clean Screen Xcel<sup>™</sup>, UCT Bristol, PA, USA) and eluted with 0.78 mL of Acetonitrile, 200 µL of 2-Propanol, and 20 µL of Ammonia. After the elution, the sample was dried under N2 gas and dissolved in 100 µL of 50% Methanol / 50% DI Water / 0.1% Formic Acid. Before injection, the sample was filtered through a 0.45 µm Nylon Syringe Filter (MicroSolv Tech Corp.).

**Note:** Opioids have been used as pain relieving drugs but are also abused as drugs producing feelings of euphoria. They are highly addictive. Nearly three out of four drug overdoses involve pain killers, so the screening of this class of drugs in urine or plasma samples is very important.



Attachment

No 312 Morphine and Hydromorphone in Urine pdf 0.2 Mb Download File

Printed from the Chrom Resource Center Copyright 2024, All Rights Apply **MicroSolv Technology Corporation** 9158 Industrial Blvd. NE, Leland, NC 28451 tel. (732) 380-8900, fax (910) 769-9435 Email: customers@mtc-usa.com Website: www.mtc-usa.com