


Acetaminophen Impurities Method with HPLC - AppNote

Acetaminophen - Alternate USP Method

Acetaminophen and two of its major impurities were analyzed using a simple Mobile Phase. The Peak Shapes are very good and the Repeatability was good (%RSD = 0.01).



× Peaks:

1. 4-Aminophenol 1.072 minutes
2. Acetaminophen 4.668 minutes
3. 4-nitrophenol 7.588 minutes

Method Conditions:

Column: Cogent Bidentate C18™, 4µm, 100Å

Catalog No.: 40018-75P

Dimensions: 4.6 x 75mm

Mobile Phase:

A: DI Water with 0.1% Formic Acid

B: Acetonitrile with 0.1% Formic Acid

Gradient:

Time (minutes)	%B
0	0
1	0
4	30
6	30
6.01	10

10

10

Post Time: 3 minutes

Injection Volume: 5µL

Flow Rate: 1.0mL / minute

Detection: UV

280 nm (4-aminophenol, acetaminophen)

324 nm (4-nitrophenol)

Sample Preparation: The stock solution was prepared by dissolving 1.0mg of standards in 10.00mL of the Mobile Phase (50:50 Solvent A / Solvent B). The Solution was then filtered with a 0.45µm Nylon Syringe Filter (MicroSolv Tech Corp.). The injection sample was diluted 1:10.

t₀: 0.9 minutes

Note: Acetaminophen (n-acetyl-p- aminophenol, APAP) is a non-steroid anti- inflammatory drug. There are several impurities that can be present in a final drug (depending on the synthetic route, the quality of starting materials, reagents, etc.) which can have safety implications. Analytical methods are needed to detect and identify these impurities and quantify them.



Attachment

No 249 Acetaminophen Impurities Method with HPLC pdf 0.3 Mb [Download File](#)

MicroSolv Technology Corporation

9158 Industrial Blvd. NE, Leland, NC 28451

tel. (732) 380-8900, fax (910) 769-9435

Email: customers@mtc-usa.com

Website: www.mtc-usa.com