
# MICROS

## Azithromycin Analyzed with HPLC ELSD – AppNote

### **Retention of Macrolide Antibiotic**

Azithromycin has weak UV absorbance and typical asymmetric peak profile with low Column efficiency in many HPLC-UV methods. This ELSD Method shows good retention and peak shape along with excellent sensitivity. This method is very reproducible with %RSD values less than 0.1%, as shown in the 10 injection overlay below.



Azithromycin

### **Method Conditions**

**Column:** Cogent Bidentate C8<sup>™</sup>, 4µm, 100Å **Catalog No.:** 40008-10P **Dimensions:** 4.6mm x 100mm Printed from the Chrom Resource Center Copyright 2024, All Rights Apply **MicroSolv Technology Corporation** 9158 Industrial Blvd. NE, Leland, NC 28451 tel. (732) 380-8900, fax (910) 769-9435 Email: customers@mtc-usa.com Website: www.mtc-usa.com

# MICROS

#### **Mobile Phase:**

A: Isopropanol

B: Acetonitrile / 0.1% Triethylamine (TEA) (v/v)

#### **Gradient:**

Time ( <i>minutes</i> )	%B
0	100
1	100
2	85
3	85
4	100
5	100

Flow rate: 1.0 mL/minute

Detection: ELSD (Evaporative Light Scattering Detector) Gain: 9; Temperature: 80°C;

Injection vol.: 1µL

Sample Preparation: Reference standards (1 mg/mL) in diluent of 50:50 Acetonitrile / DI Water (v/v)

**t**₀: 1.50 Minutes

**K':** 2

*Note:* Azithromycin is a semi-synthetic macrolide Antibiotic of the Azalide class. Azithromycin inhibits bacterial protein synthesis by binding to the 50S ribosomal subunit of the bacterial 70S ribosome.

Note 2: Capacity is determined using the following equation:  $k = (t_R - t_O)/t_O$ 

- $t_{R}$  = Retention Time of an Analyte Peak
- $t_o = Retention Time of non-Retained Peak$



Printed from the Chrom Resource Center Copyright 2024, All Rights Apply **MicroSolv Technology Corporation** 9158 Industrial Blvd. NE, Leland, NC 28451 tel. (732) 380-8900, fax (910) 769-9435 Email: customers@mtc-usa.com Website: www.mtc-usa.com