

## Comparison of Extracts in Jaboticaba Fruit Powder with LCMS – AppNote

### Phenolic Compounds Separated by HPLC with LCMS

Click [HERE](#) for Column Ordering Information.

This simple Analytical Method was used to determine the content of very potent compounds present in Jaboticaba extracts and quantified some its Phenolic contents in fermented/non-fermented fruit; estimated by using standard addition.

It is important to know the quantitation of compounds before and after fermentation to optimize the valuable antioxidants for health benefits. This method is designed to evaluate the content of the extracts and primarily focuses on comparative analysis of content in fermented versus non-fermented Jaboticaba fruit powder (JFP) produced from the species *Myrciaria jaboticaba* (Vell.) Quantification of Cyanidin 3-O- glucoside is shown in the figure 1 and other compounds in figure 2.




Figure 1:

Overlaid extracted ion chromatograms of Cyanidin 3-O-glucoside in

(A) fermented JFP and spiked fermented JFP and

(B) non-fermented JFP and spiked non-fermented JFP.

Additional Phenolic Compounds: Quercetin, Quercitrin, Cinnamic Acid, Myricetin, Ellagic Acid, Gallic Acid, Hydroxycinnamic Acid, Protocatechuic Acid, Rutin and Delphinidin-3-glucoside were

analyzed from Jaboticaba Fruit using HPLC ESI-TOF-MS; shown in *Figure 2* below.

Compare to Cyanidin-3-glucoside in both Positive (left column) and Negative (right column) ESI .







2




Figure 2:

Extracted ion chromatograms of representative phenolic compounds evaluated in Jaboticaba fruit samples[1].

## Method Conditions

**Column:** Cogent Phenyl Hydride™, 4µm, 100Å

**Catalog No.:** 69020-10P-2

**Dimensions:** 2.1 x 100mm

**Mobile Phase:**

A: DI Water with 0.1% Formic Acid

B: Acetonitrile with 0.1% Formic Acid

**Gradient:**

Time (minutes)	%B
0	5
1	10
12	90
14	90
16	5

**Flow rate:** 0.4mL / minute

**Detection:** ESI-POS/NEG Agilent Model 6210 MSD TOF Mass Spectrometer

**SamplePreparation:** Fruit of the cultivar Myrciaria jabotica.

**Final concentration:** 125mg / mL. Quantification of phenolic compounds by standard addition.

Compounds	Fermented JFP (mg/g)	Non-fermented JFP (mg/g)
Quercetin	0.264	0.009
Myricetin	0.029	-
Ellagic Acid	-	5.532
Cyanidin-3-O-Glucoside	0.180	0.952
Delphinidin-3-O-Glucoside	0.207	0.549

**Notes:** Jaboticaba (*Myrciaria Jaboticaba*) is a fruit native to Brazil that is popularly consumed as food or fermented beverages. The fruit is compared to grapes in the US and fresh fruit may begin to ferment 3-4 days after harvest. Alternate processes are used to extend shelf-life such as, jams, tarts wines and liquors.

[1] S. Watanabe, M. T Matyska-Pesek, J. De J Berrios, G. R Takeoka, J. J Pesek, "HPLC/ESI-TOFMS Identification and Quantification of Phenolic Compounds in Fermented/Non-Fermented Jaboticaba Fruit (*Myrciaria jaboticaba* (Vell.) O. Berg)", Int. J. Food Sci, 3, (5) (2018) 105-109.

**Attachment**

**A376 Comparison of Extracts in Jaboticaba Fruit Powder with LCMS pdf** 0.7 Mb [Download File](#)

Printed from the Chrom Resource Center

**MicroSolv Technology Corporation**

9158 Industrial Blvd. NE, Leland, NC 28451

tel. (732) 380-8900, fax (910) 769-9435

Email: [customers@mtc-usa.com](mailto:customers@mtc-usa.com)

Website: [www.mtc-usa.com](http://www.mtc-usa.com)

Date: 06-05-2024