
A Reproducible Method for a Phenylpyrazole Insecticide

A rapid, sensitive, and reproducible method has been developed for this Insecticide. The data below, (an overlay of 10 chromatograms) illustrates how the compound can be adequately retained and detected using this straightforward method.

A phenyl ring in the column stationary phase provides strategic use of π - π Interaction with the analyte making possible the use of a very simple, Mass Spec-friendly mobile phase with formic acid as an additive.



Fipronil

Method Conditions

Column: Cogent Phenyl Hydride[™], 4µm, 100Å Catalog No.: 69020-15P Dimensions: 4.6mm x 150mm Mobile Phase: 60:40 Acetonitrile / DI Water with 0.1% Formic Acid Injection vol.: 1µL Flow rate: 1.0mL / minute Detection: UV @ 254nm

Printed from the Chrom Resource Center Copyright 2024, All Rights Apply **MicroSolv Technology Corporation** 9158 Industrial Blvd. NE, Leland, NC 28451 tel. (732) 380-8900, fax (910) 769-9435 Email: customers@mtc-usa.com Website: www.mtc-usa.com



Sample Preparation: 0.01% Solution in 50:50 Acetonitrile / DI Water %RSD: <0.1% t_o: 1.5 Minutes K': 2.5

Notes: Fipronil is a Phenylpyrazole insecticide and Acaricide. It was originally used in agriculture as seed coatings, but is commonly used as flea and tick control in dogs and cats. Fipronil is believed to act as a noncompetitive blocker of GABA-gated chloride channels.

Note 2: Capacity is determined using the following equation: $k = (t_R - t_0)/t_0$

- t_{R} = Retention Time of an Analyte Peak
- *t_o* = Retention Time of non-Retained Peak



Printed from the Chrom Resource Center Copyright 2024, All Rights Apply **MicroSolv Technology Corporation** 9158 Industrial Blvd. NE, Leland, NC 28451 tel. (732) 380-8900, fax (910) 769-9435 Email: customers@mtc-usa.com Website: www.mtc-usa.com