

Nitrosamine Impurity Assay with HPLC - Extended AppNote

API or Impurity Assay Methods for Sartan Drugs


Nitrosamines are highly toxic and are suspected to be a human carcinogen. At high doses, some haven been shown to be a hepatotoxin that causes liver fibrosis and cancer in several animal species. Due to the toxic nature of Nitrosamines, these compounds must be monitored using reliable Methods.

In this Extended AppNote, we present two HPLC Methods with UV detection for the simultaneous detection of nine Nitrosamine Impurities, two solvents used in the manufacturing processes, and several medications of importance. The nine Nitrosamine Impurities (NDMA, NMor, NMEA, NPyr, NDEA, NPip, NDPA, NDBA, NDPHA) are separated from other pharmaceutical compounds of interest. (Figure 1 and 2.)


Two solvents, Dimethylformamide (DMF) and Dimethylacetamide (DMA) are commonly used in manufacturing processes of Sartan drugs. Residual amounts of these solvents are important to detect in both raw processes and final drug products. Currently, no published data exists for separation of all nine Nitrosamine Impurities and solvents: DMF and DMA.


Click Download File Below for Full Study Conditions.











Peaks:

| Peak Number | Abbreviation | Chemical name |
|-------------|--------------|---------------------------|
| 1 | NDMA | N-Nitrosodimethylamine |
| 2 | NMor | N-Nitrosomorpholine |
| 3 | NMEA | N-Nitrosomethylethylamine |
| 4 | Npyr | N-Nitrosopyrrolidine |
| 5 | NDEA | N-Nitrosodiethylamine |
| 6 | NPIP | N-Nitrosopiperidine |
| 7 | NDPA | N-Nitrosodi-n-propylamine |
| 8 | NDBA | N-Nitrosodi-n-butylamine |
| 9 | NDPHA | N-Nitrosodiphenvlamine |

Method Conditions:

Column 1: Cogent RP C18[™], 3um, 100Å



Catalog No.: 68318-15P

Column 2: Cogent Bidentate C18[™], 4um, 100Å

Catalog No.: 40018-15P

Dimensions: 4.6mm x 150mm (for both Columns)

Mobile Phase:

A: DI water and 0.1% Formic Acid B: Acetonitrile + 0.1% Formic Acid

Injection vol.: 1µL

Detection: UV @ 254nm and 220nm

Sample Preparation: Nitrosamine standard solution and Sartan Drug mixtures prepared as 1.0 mg/mL in Methanol

Note 1: Angiotensin receptor blockers (ARBs) are a class of medicines that is used to treat high blood pressure (also known as hypertension). These ARBs work by blocking receptors that the hormone acts on, specifically AT1 receptors, which are found in the heart, blood vessels and kidneys. Blocking the action of angiotensin II helps to lower blood pressure and prevent damage to the heart and kidneys. These medicines typically have names that end in 'sartan'. Some of these sartan drug brands are valsartan, irbesartan, candesartan, losartan and olmesartan.



Attachment

Nitrosamine Impurity Assay with HPLC - AppNote Download File

MicroSolv Technology Corporation

9158 Industrial Blvd. NE, Leland, NC 28451 tel. (732) 380-8900, fax (910) 769-9435

Email: customers@mtc-usa.com

Website: www.mtc-usa.com