


## Seven Nitroaromatic and Nitroamine Explosives Analyzed with HPLC - AppNote

## A Reproducible Method for Detection of a Nitroaromatics

Click HERE for Column Ordering Information.

The EPA Standard Test Method (8330B) for the analysis of Seven Nitroaromatic and Nitroamine Explosives in Soil by HPLC requires the use of two Reversed-Phase Columns, C18 and CN in series. These Explosive compounds are notorious for separation difficulty due to the similarities in structure.

This Method has been developed for all seven of the Nitroaromatic Explosives and is rapid, sensitive, and Reproducible .



## Peaks:

- 1. Hexahydro-1,3,5-trinitro-1,3,5-triazine (HTNT)
- 2. Octahydro-1,3,5,7-tetranitro-1,3,5,7-tetrazocine (OTNT)
  - 3. m-Dinitrobenzene (DMB)
  - 4. 2-Amino-4,6-dinitrotoluene (ADNT)
    - 5. 1,3,5-Trinitrobenzene (TNB)
      - 6. 2,4-Dinitrotoluene (DNT)
    - 7. 2,4,6-Trinitrotoluene (TNT)



## **Method Conditions**

Column: Cogent Phenyl Hydride™, 4µm, 100Å

Catalog No.: 69020-15P

Dimensions: 4.6mm x 150mm

**Mobile Phase**:

A: DI Water with 0.1% Formic Acid (v/v)
B: Acetonitrile with 0.1% Formic Acid (v/v)

**Gradient**:

Time (minutes)	%B
0	25
10	65
11	25
15	25

Injection vol.: 1µL

Flow rate: 1.0mL / minute Detection: UV @ 254nm

Sample Preparation: EPA Method 8330 Explosives Mixture each analyte @ 100 ug/mL Standard Solution in

Acetonitrile %RSD: <0.2%

Notes: These solid compounds do not occur naturally in the environment. The commonly known compound, TNT, is an explosive used in military shells, bombs, and grenades, in industrial uses, and in underwater blasting. 2,4,6-Trinitrotoluene production in the United States occurs solely at military arsenals.





Printed from the Chrom Resource Center

MicroSolv Technology Corporation

9158 Industrial Blvd. NE, Leland, NC 28451 tel. (732) 380-8900, fax (910) 769-9435

Email: customers@mtc-usa.com

Website: www.mtc-usa.com

Date: 05-07-2024