


Ticagrelor Analyzed with HPLC - AppNote

A Reproducible Method for Detection of this Blood Clot Preventative

Click **HERE** for Column Ordering Information.

A rapid, sensitive, and Reproducible Method has been developed for analysis of Ticagrelor. The data below, (overlay of 5 chromatograms) illustrates how the compound can be adequately Retained and detected using a simple Gradient in Reversed Phase HPLC. The Method demonstrates good Peak Shape and run-to-run Precision with RSD values less than 0.3%.

A Phenyl ring in the Column Stationary Phase provides beneficial π - π Interaction with the Analyte making possible the use of a very simple, Mass Spec friendly Mobile Phase with Formic Acid as an additive.



5 Injections of Ticagrelor Overlay

Printed from the Chrom Resource Center Copyright 2024, All Rights Apply **MicroSolv Technology Corporation** 9158 Industrial Blvd. NE, Leland, NC 28451

tel. (732) 380-8900, fax (910) 769-9435

Email: customers@mtc-usa.com

Website: www.mtc-usa.com



Method Conditions

Column: Cogent Phenyl Hydride[™], 4μm, 100Å

Catalog No.: 69020-10P-2

Dimensions: 2.1mm x 100mm

Mobile Phase:

A: DI Water with 0.1% Formic Acid (v/v)
B: Acetonitrile with 0.1% Formic Acid (v/v)

Gradient:

Time (minutes)	%B
0	40
1	40
5	85
6	85
7	40
10	40

Injection vol.: 1µL

Flow rate: 0.2mL / minute Detection: UV @ 254nm

Sample Preparation: 2.0mg / mL Ticagrelor standard solution in Methanol.

Notes: Ticagrelor is an oral prescription indicated for treatment of acute ischemic stroke, coronary artery disease and acute coronary syndrome.



Printed from the Chrom Resource Center Copyright 2024, All Rights Apply

MicroSolv Technology Corporation

9158 Industrial Blvd. NE, Leland, NC 28451 tel. (732) 380-8900, fax (910) 769-9435

Email: customers@mtc-usa.com

Website: www.mtc-usa.com