



## Docusate Sodium Analyzed by HPLC - AppNote

### A Simple Gradient with Good Separation

The official USP assay method for Docusate Sodium tablets calls for a Mobile Phase of 50% 7mM Ammonium Acetate / 50% Acetonitrile at 40°C. However, retention for the API and the System Suitability compound Methyl Paraben were found to be low using these conditions.

This Method not only exceeds the System Suitability resolution, but has both compounds adequately retained in order to allow for separation from other peaks that may be present in the sample. This Method is suitable for analysis of Docusate in a variety of matrices.



**Peaks:**

1. Methyl Paraben
2. Docusate Sodium

### Method Conditions

**Column:** Cogent Bidentate C18™, 4µm, 100Å

**Catalog No.:** 40018-75P

**Dimensions:** 4.6 x 75mm

**Mobile Phase:**

A: DI Water / 10 mM Ammonium Acetate

B: 95% Acetonitrile / 5% Solvent A (v/v)

Time (minutes)	%B
0	20

1	20
6	80
7	20

**Temperature:** 40°C

**Post Time:** 3 minutes

**Injection vol.:** 10µL

**Flow rate:** 1.0mL / minute

**Detection:** UV @ 210nm

**Sample Preparation:** 1.0mg / mL Docusate Sodium and 0.01mg / mL Methyl Paraben USP reference standards in diluent of 50:50 Solvent A / Solvent B. Peak identities were confirmed with individual standards.

**to:** 0.9 minutes

**Note:** Docusate is used in many laxative formulations. It is also an emulsifying, wetting, and dispersing agent. It was one of the components in the oil dispersant Corexit® used to help clean the Deepwater Horizon oil spill of 2010.



## Attachment

**No 213 Docusate Sodium Analyzed by HPLC pdf** 0.3 Mb [Download File](#)

Printed from the Chrom Resource Center

Copyright 2025, All Rights Apply

**MicroSolv Technology Corporation**

9158 Industrial Blvd. NE, Leland, NC 28451

Tel: (732) 380-8900

Fax: (910) 769-9435

Email: [customers@mtc-usa.com](mailto:customers@mtc-usa.com)

Website: [www.mtc-usa.com](http://www.mtc-usa.com)