



# Method transfer for cyclobenzaprine - AppNote

# Increased Efficiency: 4 µm to 2.2 µm

The two chromatograms compare the retention and efficiency of a Cyclobenzaprine peak using two types of Cogent Diamond Hydride columns. Figure A uses a near UHPLC 2.2 $\mu$ m phase while Figure B uses a standard 4 $\mu$ m particle size column. The results show consistent retention between the two phases (about 5% difference).

These method conditions can readily be transferred from the 4um column to the 2.2um; the advantage of the  $2.0^{\text{TM}}$  phase is the significantly higher efficiency.





#### Peak:

Cyclobenzaprine

## **Method Conditions**

**Column:** Cogent Diamond Hydride™, 2.2µm, 120Å

**Catalog No.:** <u>70200-05P-2</u> **Dimensions:** 2.1 x 50mm

**Mobile Phase:** 

A: 90% DI Water / 10% Acetonitrile / 0.1% Formic Acid (v/v)

B: Acetonitrile / 0.1% Formic Acid (v/v)

#### **Gradient:**

Time ( <i>Minutes</i> )	%B
0	90
0.5	90
8	40
9	40

**Post Time:** 3 minutes

Flow rate: 0.3mL / minute Detection: UV @ 230 nm

Injection vol.: 1µL Sample Preparation:

10mg strength Cyclobenzaprine tablet was ground and added to a 50 mL volumetric flask. A diluent of 80/20/0.1 acetonitrile / DI water / formic acid was added and the flask was sonicated for 30 minutes. It was then diluted to mark, mixed, and filtered with a  $0.45 \mu m$  nylon syringe filter (MicroSolv Tech Corp)

**Note:** Cyclobenzaprine is used to relieve muscle spasms and accompanying acute pain caused by various musculoskeletal maladies. Brand names include Amrix®, Flexeril®, and Fexmid®.



## **Attachment**

## Method Transfer Cyclobenzaprine Download File

Printed from the Chrom Resource Center Copyright 2025, All Rights Apply **MicroSolv Technology Corporation** 9158 Industrial Blvd. NE, Leland, NC 28451 Tel: (732) 380-8900

Fax: (910) 769-9435

Email: customers@mtc-usa.com

Website: www.mtc-usa.com